




asym_atl1




asym_atr1








asym_atl2








asym_atr2








reg_asym_atl1








reg_asym_atr1








reg_asym_atl2








reg_asym_atr2








asym_atl1








asym_atr1








asym_atl2








asym_atr2








reg_asym_atl1








reg_asym_atr1








reg_asym_atl2








reg_asym_atr2








asym_atl1








asym_atr1



asym_atl2



asym_atr2



The figure consists of five scatter plots arranged horizontally, each showing the distribution of differences between two beam position monitors (BPMs). The y-axis for all plots ranges from -15000 to 15000. The x-axis for each plot represents the difference in micrometers (um).

- diff_bpm1X**: The x-axis ranges from -100 to 100. The distribution is a broad, roughly circular cloud of points centered at (0,0).
- diff_bpm4aY**: The x-axis ranges from -300 to 400. The distribution is a narrow, vertically elongated ellipse centered at (0,0).
- diff_bpm4eX**: The x-axis ranges from -60 to 80. The distribution is a broad, roughly circular cloud of points centered at (0,0).
- diff_bpm4eY**: The x-axis ranges from -200 to 400. The distribution is a narrow, vertically elongated ellipse centered at (0,0).
- diff_bpm12X**: The x-axis ranges from -150 to 150. The distribution is a broad, roughly circular cloud of points centered at (0,0).