




asym_atl1




asym_atr1




asym_atl2




asym_atr2




reg_asym_atl1




reg_asym_atr1








reg_asym_atl2








reg_asym_atr2








asym_atl1








asym_atr1








asym_atl2








asym_atr2








reg_asym_atl1








reg_asym_atr1








reg_asym_atl2








reg_asym_atr2








asym_atl1








asym_atr1



asym_atl2



asym_atr2



The figure consists of five scatter plots arranged horizontally, each showing the distribution of differences between two beam position monitors (BPMs). The plots are titled as follows:

- diff_bpm1X**: The y-axis is labeled 'diff_bpm1X' and ranges from -20000 to 20000. The x-axis is labeled 'diff_bpm1X/um' and ranges from -100 to 100.
- diff_bpm4aY**: The y-axis is labeled 'diff_bpm4aY' and ranges from -20000 to 20000. The x-axis is labeled 'diff_bpm4aY/um' and ranges from -50 to 40.
- diff_bpm4eX**: The y-axis is labeled 'diff_bpm4eX' and ranges from -20000 to 20000. The x-axis is labeled 'diff_bpm4eX/um' and ranges from -100 to 100.
- diff_bpm4eY**: The y-axis is labeled 'diff_bpm4eY' and ranges from -20000 to 20000. The x-axis is labeled 'diff_bpm4eY/um' and ranges from -40 to 30.
- diff_bpm12X**: The y-axis is labeled 'diff_bpm12X' and ranges from -20000 to 20000. The x-axis is labeled 'diff_bpm12X/um' and ranges from -200 to 300.

Each plot shows a dense, roughly circular cloud of points centered around the origin (0,0), indicating a Gaussian-like distribution of the differences. The axes are labeled with units of micrometers (um) for the x-axis and no units for the y-axis.