




asym_atl1




asym_atr1




asym_atl2




asym_atr2




reg_asym_atl1




reg_asym_atr1








reg_asym_atl2








reg_asym_atr2








asym_atl1








asym_atr1








asym_atl2








asym_atr2








reg_asym_atl1








reg_asym_atr1








reg_asym_atl2








reg_asym_atr2








asym_atl1








asym_atr1



asym_atl2



asym_atr2



The figure consists of five scatter plots arranged horizontally, each showing the distribution of differences between two beam position monitors (BPMs). The plots are titled as follows:

- diff_bpm1X**: The x-axis is labeled `diff_bpm1X/um` and ranges from -40 to 40. The y-axis ranges from -10000 to 10000.
- diff_bpm4aY**: The x-axis is labeled `diff_bpm4aY/um` and ranges from -30 to 30. The y-axis ranges from -10000 to 10000.
- diff_bpm4eX**: The x-axis is labeled `diff_bpm4eX/um` and ranges from -40 to 40. The y-axis ranges from -10000 to 10000.
- diff_bpm4eY**: The x-axis is labeled `diff_bpm4eY/um` and ranges from -30 to 30. The y-axis ranges from -10000 to 10000.
- diff_bpm12X**: The x-axis is labeled `diff_bpm12X/um` and ranges from -150 to 200. The y-axis ranges from -10000 to 10000.

All plots show a dense, roughly circular cloud of points centered around the origin (0,0), indicating that the differences between the BPMs are generally small and centered around zero.