




asym\_atl1




asym\_atr1




asym\_atl2




asym\_atr2




reg\_asym\_atl1




reg\_asym\_atr1








reg\_asym\_atl2







reg\_asym\_atr2








asym\_atl1








asym\_atr1








asym\_atl2








asym\_atr2








reg\_asym\_atl1








reg\_asym\_atr1








reg\_asym\_atl2








reg\_asym\_atr2








asym\_atl1



asym\_atr1



asym\_atl2



asym\_atr2








Figure 1 displays five scatter plots showing the distribution of difference coordinates for different BPMs. The plots are titled: **diff\_bpm1X**, **diff\_bpm4aY**, **diff\_bpm4eX**, **diff\_bpm4eY**, and **diff\_bpm12X**. Each plot has a y-axis ranging from -10000 to 10000. The x-axis ranges are: **diff\_bpm1X/um** (-100 to 100), **diff\_bpm4aY/um** (-30 to 30), **diff\_bpm4eX/um** (-100 to 100), **diff\_bpm4eY/um** (-40 to 30), and **diff\_bpm12X/um** (-200 to 200). The distributions are centered around (0,0) and show varying degrees of elongation and spread.

The figure consists of five scatter plots arranged horizontally, each showing the difference between regression and postman results for a specific BPM. The plots are titled: **diff\_bpm1X**, **diff\_bpm4aY**, **diff\_bpm4eX**, **diff\_bpm4eY**, and **diff\_bpm12X**. Each plot has a y-axis labeled 'diff\_bpm' and an x-axis labeled 'diff\_bpm'. The y-axis ranges from -10000 to 10000, and the x-axis ranges from -100 to 100. The plots show a dense cloud of points centered around zero, indicating high agreement between the two methods.