




asym_atl1




asym_atr1




asym_atl2




asym_atr2




reg_asym_atl1




reg_asym_atr1








reg_asym_atl2








reg_asym_atr2








asym_atl1








asym_atr1








asym_atl2








asym_atr2








reg_asym_atl1








reg_asym_atr1








reg_asym_atl2








reg_asym_atr2








asym_atl1








asym_atr1



asym_atl2



asym_atr2



The figure consists of five scatter plots arranged horizontally, each showing the distribution of differences between a specific BPM and BPM4a. The plots are titled as follows:

- diff_bpm1X**: The x-axis is labeled $\text{diff_bpm1X}/\mu\text{m}$ and ranges from -100 to 150. The y-axis is labeled $\text{diff_bpm1X}/\mu\text{m}$ and ranges from -15000 to 15000.
- diff_bpm4aY**: The x-axis is labeled $\text{diff_bpm4aY}/\mu\text{m}$ and ranges from -30 to 30. The y-axis is labeled $\text{diff_bpm4aY}/\mu\text{m}$ and ranges from -15000 to 15000.
- diff_bpm4eX**: The x-axis is labeled $\text{diff_bpm4eX}/\mu\text{m}$ and ranges from -150 to 150. The y-axis is labeled $\text{diff_bpm4eX}/\mu\text{m}$ and ranges from -15000 to 15000.
- diff_bpm4eY**: The x-axis is labeled $\text{diff_bpm4eY}/\mu\text{m}$ and ranges from -30 to 20. The y-axis is labeled $\text{diff_bpm4eY}/\mu\text{m}$ and ranges from -15000 to 15000.
- diff_bpm12X**: The x-axis is labeled $\text{diff_bpm12X}/\mu\text{m}$ and ranges from -200 to 200. The y-axis is labeled $\text{diff_bpm12X}/\mu\text{m}$ and ranges from -15000 to 15000.

All plots show a dense, roughly circular cloud of points centered at the origin (0,0), indicating that the differences are generally small and centered around zero.

The figure displays five scatter plots, each representing the distribution of a specific difference variable. The plots are arranged horizontally. Each plot has a title above it and axes labeled with the variable name and units (um). The distributions are roughly circular and centered at (0,0).

- diff_bpm1X**: The x-axis ranges from -100 to 150, and the y-axis ranges from -10000 to 10000.
- diff_bpm4aY**: The x-axis ranges from -30 to 30, and the y-axis ranges from -10000 to 10000.
- diff_bpm4eX**: The x-axis ranges from -150 to 150, and the y-axis ranges from -10000 to 10000.
- diff_bpm4eY**: The x-axis ranges from -30 to 20, and the y-axis ranges from -10000 to 10000.
- diff_bpm12X**: The x-axis ranges from -200 to 200, and the y-axis ranges from -10000 to 10000.