




asym_usl




asym_dsl




asym_usr




asym_dsr




reg_asym_usl




reg_asym_dsl








reg_asym_usr








reg_asym_dsr








asym_usl








asym_dsl








asym_usr








asym_dsr








reg_asym_usl








reg_asym_dsl








reg_asym_usr








reg_asym_dsr








asym_usl








asym_dsl



asym_usr



asym_dsr



The figure displays five scatter plots comparing the difference between BPM4 and BPM5 for various parameters. The plots are arranged in a single row, each with a title above it. The y-axis for all plots ranges from -500 to 1500. The x-axis for each plot represents the difference between the two models for a specific parameter.

- diff_bpm4aX**: The x-axis ranges from -20 to 5. The data points are centered around 0.
- diff_bpm4aY**: The x-axis ranges from -10 to 30. The data points are centered around 0.
- diff_bpm4eX**: The x-axis ranges from -15 to 50. The data points are centered around 0.
- diff_bpm4eY**: The x-axis ranges from -40 to 40. The data points are centered around 0.
- diff_bpmE**: The x-axis ranges from -40 to 80. The data points are centered around 0.

Each plot shows a dense, roughly elliptical cloud of points, indicating that the differences between BPM4 and BPM5 are generally small and centered around zero for these parameters.