




asym_atl1




asym_atr1




asym_atl2




asym_atr2




reg_asym_atl1




reg_asym_atr1








reg_asym_atl2








reg_asym_atr2








asym_atl1








asym_atr1








asym_atl2








asym_atr2








reg_asym_atl1








reg_asym_atr1








reg_asym_atl2








reg_asym_atr2








asym_atl1








asym_atr1



asym_atl2



asym_atr2



The figure consists of five scatter plots arranged horizontally, each showing the difference between two models for a specific component. The y-axis for all plots is labeled 'diff_bpm4a' and ranges from -2000 to 2000. The x-axis for all plots is labeled 'diff_bpm4e' and ranges from -40 to 40. The plots are titled as follows:

- diff_bpm4aX**: Shows the difference for the X component. The x-axis ranges from -8 to 12.
- diff_bpm4aY**: Shows the difference for the Y component. The x-axis ranges from -40 to 40.
- diff_bpm4eX**: Shows the difference for the X component. The x-axis ranges from -60 to 80.
- diff_bpm4eY**: Shows the difference for the Y component. The x-axis ranges from -20 to 20.
- diff_bpmE**: Shows the difference for the E component. The x-axis ranges from -40 to 40.

In all plots, the data points are tightly clustered around the zero line on the x-axis, indicating a high level of agreement between the two models for all components.