




asym_atl1




asym_atr1




asym_atl2




asym_atr2




reg_asym_atl1




reg_asym_atr1








reg_asym_atl2








reg_asym_atr2








asym_atl1








asym_atr1








asym_atl2








asym_atr2








reg_asym_atl1








reg_asym_atr1



reg_asym_atl2








reg_asym_atr2








run4903_000_regression_reg_at_vs_diff_bpm-fit_postpan.png






asym_atl1








asym_atr1



asym_atl2



asym_atr2



The figure consists of five scatter plots arranged horizontally, each showing the difference between two methods for a specific coordinate. The y-axis for all plots is labeled 'diff_bpm4a' and ranges from -40,000 to 40,000. The x-axis for all plots is labeled 'diff_bpm4e' and ranges from -40,000 to 40,000. The plots are titled as follows:

- diff_bpm4aX**: Shows the difference for the X coordinate. The x-axis ranges from -6 to 6.
- diff_bpm4aY**: Shows the difference for the Y coordinate. The x-axis ranges from -10 to 10.
- diff_bpm4eX**: Shows the difference for the X coordinate. The x-axis ranges from -30 to 30.
- diff_bpm4eY**: Shows the difference for the Y coordinate. The x-axis ranges from -10 to 10.
- diff_bpmE**: Shows the difference for the E coordinate. The x-axis ranges from -40 to 40.

All plots show a dense cloud of points centered around zero, indicating high agreement between the two methods.